

Why don't patients attend their appointments? Maintaining engagement with psychiatric services

Alex J. Mitchell & Thomas Selmes

Abstract Patients miss about 20% of scheduled appointments for mental health treatment, almost twice the rate in other medical specialties. Up to 50% of patients who miss appointments drop out of scheduled care. Many who miss appointments because of slips and lapses later rearrange their appointments without adverse consequences. Those that do not are at risk of further deterioration, relapse and hospital readmission. Predictors of non-attendance are complex and linked with the predictors of missed medication. Service barriers and administrative errors are common but are often overlooked in the absence of feedback from patients. Of prime importance are the therapeutic alliance and degree of 'helpfulness' of the clinician but again these are rarely measured routinely. Useful markers of engagement include patient-rated trust, satisfaction and degree of perceived participation in treatment decisions. Much can be done to improve attendance in most services. Simple measures such as offering prompt, convenient appointments, offering reminders and augmenting with telephone contact have a reasonable evidence base. Scales to assess therapeutic alliance are now available. Complex interventions need to be evaluated carefully in order that the overall benefits outweigh costs. We suggest that clinicians consider accessibility, discharge policies and patient feedback when examining local rates of non-attendance.

This is the second of two articles by Mitchell & Selmes in APT on patient engagement and retention in treatment. The first, which addressed non-adherence to medication, appears in the previous issue of the journal (Mitchell & Selmes, 2007a).

The financial cost of missed appointments in the National Health Service (NHS) has been estimated at £360 million per year (Stone *et al*, 1999), most of this accounted for by non-attendance in primary care and hospital out-patient clinics. The purpose of this article is to review the extent and predictors of non-attendance in people with mental disorders presenting in primary and secondary care. Although one approach may be to fundamentally question the merits of out-patient clinics (Killaspy, 2006), our approach is to examine essential predictors and solutions within current models of care. In considering the classification of missed appointments it is important to distinguish between those who do not attend their first appointment and those who do

not attend follow-up appointments. Unfortunately, in reviewing the literature few studies uphold this distinction. It is also useful to separate individuals who occasionally miss appointments (partial non-attendance) and those who disengage from any follow-up (often called treatment drop-outs). To put non-attendance in context, patients may experience difficulty following medical advice at each stage of their patient journey (Fig. 1).

The extent of non-attendance in psychiatry (and perhaps its impact on the patient) may be significantly greater than in other medical specialties. In 2002–2003, 19.1% of psychiatry out-patient appointments were missed in England, compared with an NHS-wide figure of 11.7% (Department of Health, 2003). Of note, the Commission for Healthcare Improvement (now the Healthcare Commission) showed that non-attendance rates varied more than fivefold across 83 mental health trusts in England (data collated from <http://www.chi.nhs.uk/ratings>). These national figures have been replicated in numerous small studies based in psychiatry out-patient clinics, which have reported non-attendance

[†]For a commentary on this article see pp. 435–437, this issue.

Alex Mitchell is Consultant and Honorary Senior Lecturer in Liaison Psychiatry at University Hospitals Leicester (Department of Liaison Psychiatry, Brandon Unit, Leicester General Hospital, Gwendolen Road, Leicester LE5 4PW, UK. Email: alex.mitchell@leicspart.nhs.uk) and author of the prize-winning book *Neuropsychiatry and Behavioural Neurology Explained: Diseases, Diagnosis, and Management* (2003). Thomas Selmes is a senior house officer on the York psychiatry scheme and was previously a medical student at Leicester University.

rates of 15.6–28% when examined as a proportion of all offered appointments (Carpenter *et al*, 1981; Matas *et al*, 1992; Dobscha *et al*, 1999; Killaspy *et al*, 2000). Rates of initial non-attendance are usually higher than follow-up non-attendance, although this pattern may not hold for certain specialties (Mitchell & Selmes, 2007b). The frequency of partial non-attendance is higher than the rate of full discontinuation (drop-out), echoing the findings of studies of missed medication (Mitchell & Selmes, 2007a). Regarding partial non-attendance, Akerblad *et al* (2003) in Sweden found that among 1031 patients with depression treated in primary care over a 6 month period only 54.6% of all scheduled appointments were kept. In the largest study of full disengagement, Young *et al* (2000) found that 31% of community psychiatric patients were lost to follow-up over the course of 1 year. Over the course of 2 years the documented attrition rate for unscheduled drop-out may be as high as 50% (Percudani *et al*, 2002).

Consequences of non-attendance in psychiatry

Non-attendance as a predictor of drop-out, relapse and readmission

An important question is how many of those who miss an appointment later reattend and resume treatment, and how many drop out of care permanently. Early non-attendance increases the risk of further non-attendance (Carpenter *et al*, 1981), but two prospective studies suggest that many patients who occasionally miss appointments subsequently reattend. Sparr *et al* (1993) found that over a 3-month period 71.1% of missed appointments were rescheduled spontaneously by the patient (mostly within 2 weeks) and only about 25% resulted in disengagement from services. Pang *et al* (1996) traced 258 patients 6 months after their non-attendance at a follow-up clinic: 50% eventually returned but 50% dropped out of treatment. There is also good evidence that missed appointments, and unexpected disengagement in particular, can signal deteriorating mental health. In a prospective study of 365 out-patients receiving mental healthcare, Killaspy *et al* (2000) found that follow-up patients who missed an appointment were more unwell and more functionally impaired than those who attended. In a comprehensive study, Nelson *et al* (2000) collected data on 3113 psychiatric admissions in eight south-eastern US states and calculated rehospitalisation rates after discharge. On the basis of the 365-day rehospitalisation rate, patients who kept a follow-up appointment




Fig. 1 Stages of difficulty in following medical advice.

had a 1 in 10 chance of being rehospitalised; for patients who did not keep (or were not offered) an appointment, the chances were 1 in 4. It is possible that the complications of non-attendance differ for new and ongoing patients. Killaspy *et al* (2000) interviewed 57 new patients and 167 follow-up patients who did and did not attend general adult psychiatric out-patient clinic appointments over 12 months. Follow-up patients who did not attend had a 33% chance of being admitted in the next 12 months, compared with a 20% chance for the group that did attend. Also, the new patients who failed to attend were less likely to have agreed to the referral and less likely to have had it adequately explained to them.

A related concern is that attendance rates in those recently seen in emergency departments for self-harm who are offered follow-up are low (Mitchell, *et al*, 2006). Indeed, the Confidential Inquiry into Homicides and Suicides by Mentally Ill People (Royal College of Psychiatrists, 1996) found that 28% of mentally ill suicide victims had

lost contact with mental health services, 24% of suicides occurred within a month of discharge from hospital and most were before the first follow-up appointment had even taken place.

Attendance and medication adherence

Regular monitoring has been shown to lead to significant improvements in adherence to medication regimens (Fig. 2). When the US Federal Drug Administration decreased mandatory clozapine blood monitoring from weekly to every second week, there was a 10% rise in medication discontinuation (Patel *et al*, 2005). As discontinuation of medication can lead to a significant deterioration in quality of life, it can be said that clinical contacts, medication use and quality of life are interrelated. Non-attendance is particularly closely linked with medication non-adherence (Mitchell & Selmes, 2007a). Patients who discontinue medication of their own accord may be reluctant to disclose this to medical staff, and indeed cannot disclose it if not seen again. Conversely, patients who miss appointments will not receive the full benefit of medical advice and hence are less likely to make

an informed choice about their care and are more likely to act autonomously in a self-directed manner (Mitchell, 2007).

Non-attendance and medical attitudes

Missed appointments may result in increased provider frustration (Husain-Gambles *et al*, 2004), decreased levels of provider empathy and lower quality patient-provider communication (Pesata *et al*, 1999). In a primary care study, general practitioners (GPs) developed negative attitudes about patients who unexpectedly missed their appointments (Husain-Gambles *et al*, 2004). Not infrequently, favoured intervention strategies involve punishing the patient in some way for their 'offence' rather than discussing missed appointments with the patient (see also the section 'Service responses to missed appointments' below). Remarkably, despite the high risk of missed appointments in mental health services, a US survey showed that psychiatrists tended to be less active in trying to maintain contact with patients than non-physician therapists and junior doctors (Smoller *et al*, 1998).




Fig. 2 Illustration of the factors influencing adherence to medical advice.

Predictors of non-attendance

Few studies have examined the predictors of non-attendance systematically in mental health settings (Chen, 1991). In general, predictors may be divided into environmental and demographic factors, patient factors, illness factors and clinician factors (Box 1). The latter has been least investigated.

Box 1 Key predictors of non-attendance

Environmental and demographic factors

- Younger age
- Lower socio-economic status
- Not having health insurance (where health-care is not free at point of delivery)
- Poor adherence to psychotropic medications
- Homelessness
- Transport problems, distance from clinic

Patient factors

- Forgetting, oversleeping, getting the date wrong
- Being too psychiatrically unwell
- High trait anxiety
- Lower social desirability scores
- Dismissing attachment styles

Memory/cognitive problems

- Dementia

Information and health beliefs

- Poor insight into illness

Illness factors

- Personality disorder
- Substance misuse (alone or in combination with other psychiatric disorder)
- Neurotic disorders
- Diagnosis unclear or cannot be established

Clinician and referrer factors

- Poor communication between the referring practitioner and the patient
- Patient's disagreement with the referral
- Referrer's scepticism about the value of psychiatry
- Poor-quality referral letter
- Longer delay between the referral and the appointment (or between assessment and treatment)
- Early stages of treatment
- Quality of therapeutic alliance
- Non-collaborative decision-making

Environmental and demographic factors

A number of social and environmental factors have been linked to missed appointments in most medical specialties. Research indicates that patients who miss appointments tend to be younger and of lower socio-economic status (Sharp & Hamilton, 2001). Compared with attenders, such patients are more often in receipt of healthcare (incapacity) benefits and/or social benefits and are more likely to have psychosocial problems (Cosgrove, 1990). In a comprehensive study of 221 000 primary care appointments in Sheffield, there was a strong correlation ($r=0.72$) between non-attendance and the Townsend Deprivation Index Score for the patient's medical practice (Waller & Hodgkin, 2000). Kruse *et al* (2002) examined the medical records of 313 individuals with serious mental illness who were attending a US psychiatric out-patients clinic and found that 36% had missed their first appointment. The significant predictors were young age, Hispanic ethnicity, having a poor family support system, not having health insurance and also poor adherence to psychotropic medication (Kruse *et al*, 2002). In a study of 600 patients followed up after an emergency psychiatric assessment, Dobscha *et al* (1999) found that homeless people were the least likely to attend, managing only a 34% follow-up rate. Although encouragement from family and friends is helpful, people who feel pressurised into attending by their family or doctor may be less likely to keep appointments.

A number of understandable accessibility issues are common to most studies of both help-seeking and attendance. Most frequently, transportation problems are cited (Pesata *et al*, 1999), and distance travelled is another correlate (Jackson *et al*, 2006).

Patient factors

The most common single reason for non-attendance at follow-up appointments is forgetting the appointment, followed by being too psychiatrically unwell (Killaspy *et al*, 2000). The proportion of psychiatric patients accidentally (unintentionally) missing an appointment appears to be higher than in other medical specialties. Sparr *et al* (1993) examined reasons for non-attendance in 8.8% of 1620 contacts who missed scheduled appointments during a 3-month period. The most commonly stated reason for missing an appointment was patient error (forgetting, oversleeping or getting the date wrong). Gudjonsson *et al* (2004) examined non-attendance in an alcohol clinic in Reykjavik. Of 313 patients asked to attend a follow-up appointment, only 27% did so. Predictors of non-attendance were younger

age, high trait anxiety and lower social desirability scores. Psychoticism on the Eysenck Personality Questionnaire was the biggest single discriminator between those who attended. A study of the management of 3923 patients with diabetes in primary care found that those with secure attachment styles (as assessed by the Relationship Questionnaire) were significantly more likely to attend than those with dismissing attachment styles (Ciechanowski *et al*, 2006). Compton *et al* (2006) collected data from team members and medical notes on 234 consecutively discharged patients from two in-patient units in a US urban county hospital. In a logistic regression model, the four characteristics that were significant predictors of missed first appointments were involuntary legal status at discharge or leaving against medical advice, not having an established out-patient clinician, an Axis IV problem relating to the patient's primary support group (for example, death of a family member or health problems in the family) and the number of days from discharge to the follow-up appointment.

Memory problems

Memory impairment may play a role in non-attendance as well as in subsequent engagement with treatment and services. Rosen *et al* (2003) examined the role of cognition in adherence behaviour in patients with diabetes. Low scores on the Mini-Mental State Examination and 'non-Caucasian' ethnicity were associated with missed appointments. McKellar *et al* (2006) found that greater cognitive dysfunction predicted drop out from treatment in a residential substance use disorder programme. This effect is not a reflection of old age itself, as older people have lower rates of non-attendance than younger patients (Mitchell & Selmes, 2007b). A further concern with cognitive impairment is that it may affect a patient's ability to engage with a consultation or treatment programme – or indeed to subsequently recall the results of consultations. Neupert & McDonald-Miszczak (2004) studied 105 younger adults and 58 older adults who did not have a psychiatric diagnosis. Hierarchical regression results revealed that cognitive abilities and metacognitive beliefs were related to delayed recall of medication instructions. Dementia is an obvious reason for non-attendance but this has been inadequately studied (Dockery *et al*, 2001).

Information and health beliefs

Several authors have observed that patients with schizophrenia who miss appointments are more likely to have lower levels of functioning, to be

more severely unwell and also to have substance use problems (Coodin *et al*, 2004). Yet, in depression and anxiety disorders, patients with milder disorders appear less likely to attend (Issakidis & Andrews, 2004). It is probable that a bimodal (U-shaped) relationship is present, with both the mildest and the most severe disorders linked with greatest non-attendance. Severity of illness and non-attendance may be mediated by insight. In a study of 33 patients with psychosis, poor scores on all commonly used insight assessment scales predicted poor adherence to treatment (Sanz *et al*, 1998). Most significantly, a large systematic review of 103 studies into clinic attendance and medication adherence in psychosis found an overall non-adherence rate of 25.8%. Predictors of non-adherence were lack of insight, positive symptoms, younger age, male gender, substance misuse, unemployment and poor social functioning (Nose *et al*, 2003).

Illness type

Matas *et al* (1992) reviewed the notes of 874 new out-patient referrals, and found that non-attendance was associated with a diagnosis of a personality disorder or substance misuse.

Percudani *et al* (2002) found that in an Italian sample, patients with neurotic and personality disorders were more likely to drop out of treatment than those with schizophrenia. Similarly, Young *et al* (2000) found that patients with schizophrenia were more likely to remain in contact with routine mental health services in Ventura County, California. Interestingly, if the diagnosis is unclear (or cannot be established) patients are more likely to drop out of care (Melo & Guimarães, 2005).

Bell (2001) reviewed the evidence on factors influencing disengagement with treatment for bulimia nervosa. Although 28 studies were found, the only robust finding was that comorbid borderline personality disorder increased the likelihood of withdrawal from treatment. These differences are also reflected in different rates of non-attendance across services. We examined 17211 out-patient psychiatric contacts across one calendar year and found a hierarchy of non-attendance (Mitchell & Selmes, 2007a). Non-attendance rates for new patients ranged from 36.9% in alcohol services and 25.3% in drug services to 5.0% in the treatment and recovery (rehabilitation) service and 8.7% in liaison psychiatry. In most psychiatric subspecialties, the follow-up non-attendance rate was less than that for new patients. However, this pattern was reversed in child and adolescent psychiatry, drugs services, old age psychiatry and, most notably, liaison

psychiatry, where the 8.7% of new-patient non-attendance rose to 20.9% at follow-up. Finally, many studies show that substance misuse complicates attendance. For example, Sparr *et al* (1993) showed that patients with post-traumatic stress disorder and/or substance misuse were significantly more likely than others to miss appointments. In a study of 144 patients, Ford *et al* (1991) showed that those with comorbid substance misuse failed to keep 27.5% of scheduled appointments, compared with 15.8% of those with uncomplicated psychiatric disorders. In addition, the dual-diagnosis group had a drop-out rate almost three times higher than that of the single-diagnosis group.

Clinician and referrer factors

There is no doubt that the quality of service provided, as well as the rapport with the clinician, have a marked effect on attendance. Wilder *et al* (1997) suggest that 'compliance' is 'as much a function of their [patients'] interactions with psychiatric personnel and of the suitability of the recommendations as it is of personal characteristics of the patients themselves'. In fact, this effect begins with the original referrer to secondary care.

Influence of the referrer

Poor communication between the referring practitioner and the patient may increase non-attendance at an initial appointment. Indeed, patients who agree with their referral are more likely to attend than those who do not (Killaspy *et al*, 2000). General practitioner referrals to specialist care may result in a lower rate of attendance than those from other specialists (Carpenter *et al*, 1981). In a recent study (Mitchell & Selmes, 2007b), we found that the rate of both initial and subsequent missed appointments was highest following self-referral and referrals from the police/probation service and lowest from community psychiatric nurses (CPNs), social services and from within psychiatric services. Farid & Alapont (1993) found a remarkable association between non-attendance and a referrer who was sceptical about the value of psychiatry. There was also an association between the quality of the referral letter and the likelihood of attendance.

Influence of the assessor (primary clinician)

On a simplistic level, it is interesting to note that rates of missed appointments with consultants (clinical psychologists and psychiatrists) appear to be lower than those with trainees (McIvor *et al*, 2004). Rates of initial non-attendance also increase with time when there is a delay between the referral (or scheduling)

and the actual appointment (Grunebaum *et al*, 1996; Gallucci *et al*, 2005). In the same manner, rates of follow-up non-attendance increase with delay between assessment and treatment (Jackson *et al*, 2006). Several studies suggest that patients are most likely to drop out during the early phase of treatment. For example, 69% of losses to treatment occurred in the first month in one study in Brazil (Melo & Guimarães, 2005).

However, it is also useful to look at communication and interpersonal variables in more detail, utilising studies across medicine, as many clinicians will feel they have more control over communication style than scheduling delays. Perhaps the most important interpersonal variable in predicting attendance is the quality of the therapeutic alliance and, more specifically, the degree of 'helpfulness' of the health professional as perceived by the patient (Johansson & Eklund, 2006) (Box 2). That said, in some studies clinician-rated alliance is a

Box 2 Hypothetical patient's perspective of non-attendance

Anna experiences a racing heartbeat and wonders whether it is something medically serious or perhaps due to anxiety. She decides to call the GP for an appointment. The last time she had felt like this the GP had advised her to 'wait and see if things got worse', which did not seem very helpful but the symptoms did go away. The only available appointment is in 5 days' time; Anna feels that the staff are not treating her concern seriously, but accepts the appointment because she is concerned. She begins to worry that the palpitations might cause a heart attack. Then she starts to worry what kinds of tests the doctor is going to perform; she is sure any tests will be uncomfortable and worries that they might show something serious. Both make her nervous about attending. After a few days she becomes less worried and the palpitations go away. Finally, thinking that her condition must not be serious, she decides not to keep her appointment. She is not sure that the GP could have helped anyway. She does not bother to telephone and tell the clinic because they are so busy they will not even notice her absence and will probably see someone else in her place. Also it might give the doctor 10 minutes to catch up on their work (because they always seem to be in a rush to finish).

Adapted from Lacy *et al*, 2004

stronger predictor of drop-out than patient-rated therapeutic alliance (Meier *et al*, 2006). Willingness to discuss mental health issues with a doctor is also predicted largely by the perceived helpfulness of that doctor, and in turn trust in the doctor and the medical profession.

Although trust and helpfulness are rarely measured routinely, satisfaction may be a proxy measure. Global dissatisfaction with care is a strong predictor of dropping out of care in most (Pekarik, 1983; Tehrani *et al*, 1996; Young *et al*, 2000; Rossi *et al*, 2002) but not all (Killaspy *et al*, 1998) studies. However, satisfaction may be good with some aspects of care and poor with others. One study found that dissatisfaction with the skills and behaviour of health professionals and satisfaction with received treatment both predicted drop-out, suggesting that satisfaction with quality of care and satisfaction with treatment success should be separated (Rossi *et al*, 2002).

Other work suggests that satisfaction is related to a patient-centred communication style and a willingness to forge an agreed therapeutic strategy (Nakanishi *et al*, 2006). Yet patients are not always involved fully in treatment decisions. In one study, for example, only 1 in 10 patients taking clozapine remembered discussing potentially serious haematological effects (Angermeyer *et al*, 2001). The majority of patients taking antipsychotics do not feel involved in their treatment, and simply take their medication because they are asked to do so (Gray *et al*, 2005). A collaborative decision-making communication style (as reported by patients) was shown in a survey of 100 users of antidepressants to predict attendance and medication use (Bultman & Svarstad, 2000).

Conversely, bad experiences of health services have been linked with non-attendance. Gonzalez *et al* (2005) in Texas looked at adherence to appointments and medication in 95 combat veterans prescribed antidepressant medication, observed over 6 months. Appointment attendance was best in those who felt ready for treatment, perceived its benefits, had fewer negative effects from previous mental health treatment and saw their physician as collaborative. Patients who receive appointments after greater delay or who feel that the healthcare professional did not listen to, respect or understand them, are less likely to reattend (Moore *et al*, 2004).

Interventions to reduce non-attendance

Reviews are available of methods of improving engagement in psychotherapy (Ogrodniczuk *et al*,

2005) and in adult mental health and substance misuse settings (Lefforge *et al*, 2007). In addition, Macharia *et al* (1992) identified 23 randomised controlled trials in mixed settings which used attendance as the primary outcome measure. Written and telephone prompts positively influenced attendance. A more recent Cochrane review (Reda & Makhoul, 2001) found evidence that a telephone or written prompt delivered 24 h before a clinic appointment may encourage attendance. A national campaign 'Keep it or cancel it', organised by the UK government-funded Developing Patient Partnerships Agency, was accompanied by £20 million in funding for a new computerised booking system (BBC, 1999). The cost-effectiveness of enhanced engagement methods has been questioned (Sharp & Hamilton, 2001), leading to several economic cost-benefit analyses. It may be useful for the following section to consider patient's reasons for non-attendance in two roughly equal categories: avoidable reasons such as forgetting, communication problems and transport problems, and unavoidable reasons such as illness relapse and illness remission. Interventions have been mainly aimed at the so-called avoidable factors.

Reminder and memory aids

The discovery that psychiatric patients are twice as likely to cite forgetting as the cause of non-attendance than those in other specialties (Killaspy *et al*, 2000) might support interest in active reminder systems. Boswell *et al* (1983) was one of the first to use a telephone reminder before psychiatric clinic appointments for 184 patients compared with 185 treated as usual. There was a statistically significant improvement in attendance in this early randomised controlled trial. Similar positive results have since been reported for first attenders by Burgoyne *et al* (1983), MacDonald *et al* (2000) and Conduit *et al* (2004), and for both first attenders and follow-up patients by Shivack & Sullivan (1989) and for follow-up patients by Bamford *et al* (2004), the latter using both telephone and postal reminders with alcohol misusers. Telephone reminders seem to be particularly effective for elderly patients, especially those with dementia (Dockery *et al*, 2001). Messages left on an answerphone can also be successful (Conduit *et al*, 2004). McFall *et al* (2000) conducted a pilot outreach intervention for combat veterans receiving disability benefits for post-traumatic stress disorder but not receiving treatment. A letter inviting them to seek help followed by a telephone reminder led to a 23% attendance rate compared with 7% attendance for those who received just a letter. In one study of 103 new out-patients at a London psychiatric teaching

hospital, patients were asked to confirm in writing or by telephone if they still required an appointment (Baggaley, 1993). A number of automated methods of sending reminders to patients about scheduled appointments have been developed. For example, Andrade *et al* (2005) measured the effect of a memory-prompting device (Disease Management Assistance System) on adherence to treatment in people with HIV. Interestingly, adherence improved only in those with diagnosed memory impairment. Unfortunately, the expense of these systems usually proves prohibitive for the majority of patients and healthcare providers.

Interpersonal and communication issues

Simple measures such as providing patients with sufficient information about the practical aspects of their appointment (where to park, directions, etc.) and even 'orientation statements' (explaining what to expect in the consultation) can lead to improvements in attendance (Kluger & Karras, 1983). Wenning & King (1995) offered the parents of children newly referred to child psychiatry an 'orientation group meeting', in which they were informed about the service provided. This reduced non-attendance at the first (but not subsequent) clinic appointments, and also allowed parents who were unlikely to attend to be identified in advance. Patients' understanding can be increased with a simple discharge or exit interview explaining the need for follow-up and the rationale for medication. This approach appears to reduce subsequent non-attendance (Guse *et al*, 2003). Given that one of the most important interpersonal variables in predicting attendance is the degree of helpfulness of the health professional (see above) we suggest that all clinicians check whether there is anything further that can be done to help at the close of all appointments and prior to hospital discharge. Even greater insight may be gained by routinely using an instrument to check the quality of their therapeutic relationships (McGuire-Snieckus *et al*, 2007).

Service responses to missed appointments

All services must carefully consider their approach to those who miss one or more appointments (Cruz *et al*, 2001). Clearly, many patients miss a single appointment by mistake (unintentionally) and we suggest that after one missed appointment a reminder is sent to the patient with another appointment date (or at least the offer of an appointment), possibly suggesting a home visit. After a second and third consecutive unexplained miss, there is

a higher chance that the patient is unwilling or unable to attend. We suggest speaking directly to the patient, clarifying whether an appointment is still needed and, if so, finding a mutually agreed time. If a patient cannot be contacted an alternative is to ask the referrer to clarify whether the appointment is still wanted. Simply sending three written appointments for fixed out-patient slots may not be satisfactory as there is no allowance for administrative error or severe illness. An important question that arises when a patient is initially seen but then disengages is whether that service was truly meeting the individual's needs. It is not uncommon for patients to drop out of treatment because of a perceived lack of progress. For this reason we suggest attempting to ask all patients who disengage for feedback either verbally or in writing.

It is vital that services in all specialties review the ongoing needs of the patients they serve. For example, when staff at a paediatric clinic noted a particularly high non-attendance rate among Asian patients (50%) they found that common-sense steps, including avoiding appointments during Islamic religious festivals, reducing language barriers through the use of link workers, liaison with elders and education of staff, reduced this figure to 13.5% (Gatrad, 2000). Surveys of non-attending patients (as described above) show that people often view services as inaccessible (and, to a lesser extent, inflexible). Although it is uncommon for patients to cite problems with the appointment time as a reason for non-attendance, significantly more patients will attend afternoon than morning appointments (Weinerman *et al*, 2003). In a study of 118 substance-misusing young people, a community-based treatment programme that was much easier to access than conventional care reduced disengagement from 22% to 2% (Henggeler *et al*, 1996). However, flexible but complex systems risk alienating those who cannot read English, or who have intellectual disabilities or significant psychiatric morbidity (Sharp & Hamilton, 2001).

The effectiveness of providing evening and weekend appointments has not been investigated in mental health settings, but providing hospital transport or taxis is beneficial (reducing non-attendance at prenatal appointments for low-income women by 22%) (Melnikow *et al*, 1997). The innovative but controversial method of rewarding those who attend clinic and/or punishing those who do not has been successfully piloted by more than one group (Parrish *et al*, 1986; Bech, 2005). It may be more ethically acceptable to offer a nominal reward option alone ('as a thank you for letting us know you could not attend') (Pollastri *et al*, 2005).

Assertive outreach and home treatment services

A minority of patients find engagement with hospital-based services difficult or impossible. They may have a history of disengagement leading to hospital admissions. In the past two decades, dedicated assertive outreach teams have been set up in the UK with the aim of engaging with patients and reducing hospital admission. The model of assertive outreach includes long-term engagement, shared case-loads and flexibility (Sainsbury Centre for Mental Health, 2001). There is evidence that this approach is accompanied by a significant improvement in patient engagement (Wharne, 2005). Home treatment alone may also have demonstrable benefits, although not necessarily in rates of rehospitalisation (Frederick *et al*, 2002). Of course, these principles can equally be applied in routine settings and for other types of patient. It may be advantageous to engage with patients who have cognitive impairments in their own home. In a 4-year study of out-patient old age psychiatry in an urban area of the UK, home visits had a much lower 'non-attendance' rate (1.7%) than clinics (21.2%) (Anderson & Aquilina, 2002).

Conclusions

There are many reasons for missed appointments, most often slips and lapses (e.g. forgotten appointments), and in many cases patients will later reattend without adverse consequences. However, an important proportion will not reattend and deteriorate without accessing further medical help. Both missed appointments and missed medication may herald disengagement (drop-out) from care. Faced with non-attendance it is difficult for clinicians to know which category (no help needed or at risk) individual patients fall into. For this reason a policy of discharge after one or more missed appointments may not be appropriate without clarification of need. Non-attendance due to barriers to care and administrative errors is not uncommon and can often be improved (Stone *et al*, 1999). Systematic changes in appointment systems can reduce barriers to help-seeking and hence reduce non-attendance. It is important to recognise that non-attendance is not limited to those with psychotic disorders who lose insight: in fact, patients with schizophrenia may engage more successfully than other groups with mental illness (Young *et al*, 2000; Percudani *et al*, 2002).

Patient surveys reveal that most people feel anxious about seeing a doctor, especially a psychiatrist and/or when potentially receiving 'bad news' (Grunebaum *et al*, 1996). Teams can do

much to alleviate this anxiety – and hence large improvements in missed appointment rates can be achieved by relatively simple measures (Box 3).

A valid question is why many do not ring to cancel rather than missing an appointment. Preliminary data suggest that a third forget or do not consider cancelling and may be too embarrassed on the day to call to admit this, and a small but important group are too unwell and/or cannot easily ring on the day (Hamilton, 2001). Another important reason for non-attendance is problems in the therapeutic alliance. These issues are often ignored, perhaps because health professionals are uncomfortable examining their own practice. However, simple

Box 3 Simple measures to reduce non-attendance

Improving initial attendance

- Encourage referrers to explain the purpose of the referral
- Schedule the appointment as soon as possible
- Write to the patient with clear directions and explaining the mechanism of referral
- Offer the option of an afternoon appointment
- Offer the option of a community/home visit if the patient is too unwell to attend
- Consider a reminder telephone call the day before the appointment (if the patient has a telephone)

Improving follow-up attendance

- Give the patient a choice of appointment dates and/or locations
- Schedule the appointment as soon as possible
- Where possible, agree the duration of the treatment course at the start
- Work towards establishing and maintaining a good therapeutic relationship
- Involve the patient in treatment decisions

Response to missed appointments

- Contact the patient by letter or telephone
- Identify any patient-cited barriers to attending
- Confirm that the patient wishes to attend
- Affirm that the patient can still be seen without prejudice
- If possible convey hope that there is a definite prospect of improvement
- Reschedule the missed appointment as soon as possible

(Adapted from Pettinati *et al*, 2004)

measures of clinician- and patient-rated therapeutic alliance are available. High non-attendance rates are sometimes a marker that aspects of quality of care could be improved. There appears to be an important difference between psychiatry patients who miss their first appointment and those who miss a follow-up appointment. The first appointment is heavily influenced by an individual's pre-existing health beliefs and the quality of the information given by the first referrer. Attendance at follow-up appointments is a reflection both of the patient's overall satisfaction with care and the perceived need for further help (Box 2).

There appears to be a bimodal relationship between severity of illness and attendance. Low and high illness severity predict non-attendance. In other words the most common reason for not attending a first appointment is feeling better but the most serious is feeling too unwell (Cosgrove, 1990).

We suggest that all health professionals consider barriers to care and attempt to contact patients directly before discharging those who do not attend without explanation.

Declaration of interest

None.

References

- Akerblad, A. C., Bengtsson, F., Ekselius, L., *et al* (2003) Effects of an educational compliance enhancement programme and therapeutic drug monitoring on treatment adherence in depressed patients managed by general practitioners. *International Clinical Psychopharmacology*, **18**, 347–354.
- Anderson, D. N. & Aquilina, C. (2002) Domiciliary clinics. I: Effects on non-attendance. *International Journal of Geriatric Psychiatry*, **17**, 941–944.
- Andrade, A., Mcgruder, H., Wu, A., *et al* (2005) A programmable prompting device improves adherence to highly active antiretroviral therapy in HIV-infected subjects with memory impairment. *Clinical Infectious Diseases*, **41**, 875–882.
- Angermeyer, M. C., Löffler, W., Müller, P., *et al* (2001) Patients' and relatives' assessment of clozapine treatment. *Psychological Medicine*, **31**, 509–517.
- Baggaley, M. (1993) Improving the attendance for new psychiatric out-patient referrals. *Psychiatric Bulletin*, **17**, 347–348.
- Bamford, Z., Booth, P. G., McGuire, J., *et al* (2004) Maximizing patient follow-up after alcohol treatment: the effect of a threestep reminding system on response rates. *Journal of Substance Use*, **9**, 36–43.
- BBC (1999) Health: £20m boost for NHS appointments system (2 September). <http://news.bbc.co.uk/1/hi/health/435986.stm>
- Bech, M. (2005) The economics of non-attendance and the expected effect of charging a fine on non-attendees. *Health Policy*, **74**, 181–191.
- Bell, L. (2001) What predicts failure to engage in or drop out from treatment for bulimia nervosa and what implications does this have for treatment? *Clinical Psychology and Psychotherapy*, **8**, 424–435.
- Boswell, P. C., Brauzer, B., Postlethwaite, N., *et al* (1983) Improving aftercare patients' compliance with appointments through phone calls and letters. *Hospital and Community Psychiatry*, **34**, 358–360.
- Bultman, D. C. & Svarstad, B. L. (2000) Effects of physician communication style on client medication beliefs and adherence with antidepressant treatment. *Patient Education and Counselling*, **40**, 173–185.
- Burgoyne, R. W., Acosta, F. X. & Yamamoto, J. (1983) Telephone prompting to increase attendance at a psychiatric outpatient clinic. *American Journal of Psychiatry*, **140**, 345–347.
- Carpenter, P. J., Morrow, G. R., Del Gaudio, A. C., *et al* (1981) Who keeps the first outpatient appointment? *American Journal of Psychiatry*, **138**, 102–105.
- Chen, A. (1991) Noncompliance in community psychiatry: a review of clinical interventions. *Hospital and Community Psychiatry*, **42**, 282–287.
- Ciechanowski, P., Russo, J., Katon, W., *et al* (2006) Where is the patient? The association of psychosocial factors and missed primary care appointments in patients with diabetes. *General Hospital Psychiatry*, **28**, 9–17.
- Compton, M. T., Rudisch, B. E., Craw, J., *et al* (2006) Predictors of missed first appointments at community mental health centers after psychiatric hospitalization. *Psychiatric Services*, **57**, 531–537.
- Conduit, T., Byrne, S., Court, J., *et al* (2004) Non-attendance at a university-based psychology clinic: telephone appointment reminders versus no reminders. *Australian Psychologist*, **39**, 68–75.
- Coodin, S., Staley, D., Cortens, B., *et al* (2004) Patient factors associated with missed appointments in persons with schizophrenia. *Canadian Journal of Psychiatry*, **49**, 145–148.
- Cosgrove, M. P. (1990) Defaulters in general practice: reasons for default and patterns of attendance. *British Journal of General Practice*, **40**, 50–52.
- Cruz, M., Cruz, R. F. & McEldoon, W. (2001) Best practice for managing noncompliance with psychiatric appointments in community-based care. *Psychiatric Services*, **52**, 1443–1445.
- Department of Health (2003) *Hospital Activity Statistics, London*. Department of Health http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/AnnualReports/Browsable/DH_5217175
- Dobscha, S., Delucchi, K. & Young, M. (1999) Adherence with referrals for outpatient follow-up from a VA psychiatry emergency room. *Community Mental Health Journal*, **35**, 451–458.
- Dockery, F., Rajkumar, C., Chapman, C., *et al* (2001) The effect of reminder calls in reducing non-attendance rates at care of the elderly clinics. *Postgraduate Medical Journal*, **77**, 37–39.
- Farid, B. T. & Alapont, E. (1993) Patients who fail to attend their first psychiatric appointment. Non-attendance or inappropriate referral? *Journal of Mental Health*, **2**, 81–83.
- Ford, L., Snowden, L. & Walser, E. (1991) Outpatient mental health and the dual-diagnosis patient: utilization of services and community adjustment. *Evaluation and Program Planning*, **14**, 291–298.
- Frederick, S., Caldwell, K. & Rubio, D. M. (2002) Home-based treatment, rates of ambulatory follow-up, and psychiatric rehospitalization in a Medicaid managed care population. *Journal of Behavioral Health Services and Research*, **29**, 466–475.
- Gallucci, G., Swartz, W. & Hackerman, F. (2005) Impact of the wait for an initial appointment on the rate of kept appointments at a mental health center. *Psychiatric Services*, **56**, 344–346.
- Gatrad, A. R. (2000) A completed audit to reduce hospital outpatients non-attendance rates. *Archives of Disease in Childhood*, **82**, 59–61.
- Gonzalez, J., Williams, J., Noel, P. & Lee, S. (2005) Adherence to mental health treatment in a primary care clinic. *Journal of the American Board of Family Practice*, **18**, 87–96.
- Gray, R., Rofail, D., Allen, J., *et al* (2005) A survey of patient satisfaction with and subjective experiences of treatment with antipsychotic medication. *Journal of Advanced Nursing*, **52**, 31–37.
- Grunebaum, M., Luber, P., Callahan, M., *et al* (1996) Predictors of missed appointments for psychiatric consultations in a primary care clinic. *Psychiatric Services*, **47**, 848–852.

- Gudjonsson, G., Hannesdottir, K., Agustsson, T., et al (2004) Personality predictors of the failure of alcoholics to come for follow-up assessment. *Personality and Individual Differences*, **37**, 805–813.
- Guse, C. E., Richardson, L., Carle, M., et al (2003) The effect of exit-interview patient education on no-show rates at a family practice residency clinic. *Journal of the American Board of Family Practice*, **16**, 399–404.
- Hamilton, W. (2001) *General Practice Non-Attendance. A Questionnaire Survey*. NHS National Research Register.
- Henggeler, S. W., Pickrel, S. G., Brondino, M. J., et al (1996) Eliminating (almost) treatment dropout of substance abusing or dependent delinquents through home-based multisystemic therapy. *American Journal of Psychiatry*, **153**, 427–428.
- Husain-Gambles, M., Neal, R. D., Dempsey, O., et al (2004) Missed appointments in primary care: questionnaire and focus group study of health professionals. *British Journal of General Practice*, **54**, 108–113.
- Issakidis, C. & Andrews, G. (2004) Pretreatment attrition and dropout in an outpatient clinic for anxiety disorders. *Acta Psychiatrica Scandinavica*, **109**, 426–433.
- Jackson, K., Booth, P., McGuire, J., et al (2006) Predictors of starting and remaining in treatment at a specialist alcohol clinic. *Journal of Substance Use*, **11**, 89–100.
- Johansson, H. & Eklund, M. (2006) Helping alliance and early dropout from psychiatric out-patient care: the influence of patient factors. *Social Psychiatry and Psychiatric Epidemiology*, **41**, 140–147.
- Killaspy, H. (2006) Psychiatric out-patient services: origins and future. *Advances in Psychiatric Treatment*, **12**, 309–319.
- Killaspy, H., Gledhill, J. & Banerjee, S. (1998) Satisfaction of attenders and non-attenders with their treatments at psychiatric out-patient clinics. *Psychiatric Bulletin*, **22**, 612–615.
- Killaspy, H., Banerjee, S., King, M., et al (2000) Prospective controlled study of psychiatric out-patient non-attendance. Characteristics and outcome. *British Journal of Psychiatry*, **176**, 160–165.
- Kluger, M. & Karras, A. (1983) Strategies for reducing missed initial appointments in a community mental health centre. *Community Mental Health Journal*, **19**, 137–143.
- Kruse, G. R., Rohland, B. M. & Wu, X. (2002) Factors associated with missed first appointments at a psychiatric clinic. *Psychiatric Services*, **53**, 1173–1176.
- Lacy, N. L., Paulman, A., Reuter, M. D., et al (2004) Why we don't come. Patient perceptions on no-shows. *Annals of Family Medicine*, **2**, 541–545.
- Lefforge, N. L., Donohue, B. & Strada, M. J. (2007) Improving session attendance in mental health and substance abuse settings: a review of controlled studies. *Behavior Therapy*, **38**, 1–22.
- MacDonald, J., Brown, N. & Ellis, P. (2000) Using telephone prompts to improve initial attendance at a community mental health center. *Psychiatric Services*, **51**, 812–814.
- Macharia, W. M., Leon, G., Rowe, B. H., et al (1992) An overview of interventions to improve compliance with appointment keeping for keeping medical services. *JAMA*, **267**, 1813–1817.
- Matas, M., Staley, D. & Griffin, W. (1992) A profile of the noncompliant patient: a thirty-month review of outpatient psychiatric referrals. *General Hospital Psychiatry*, **14**, 124–130.
- McFall, M., Malte, C., Fontana, A., et al (2000) Effects of an outreach intervention on use of mental health services by veterans with posttraumatic stress disorder. *Psychiatric Services*, **51**, 369–374.
- McGuire-Snieckus, R., McCabe, R., Catty, J., et al (2007) A new scale to assess the therapeutic relationship in community mental health care: STAR. *Psychological Medicine*, **37**, 85–95.
- McIvor, R., Ek, E. & Carson, J. (2004) Non-attendance rates among patients attending different grades of psychiatrist and a clinical psychologist within a community mental health clinic. *Psychiatric Bulletin*, **28**, 5–7.
- McKellar, J., Kelly, J., Harris, A., et al (2006) Pretreatment and during treatment risk factors for dropout among patients with substance use disorders. *Addictive Behaviors*, **31**, 450–460.
- Meier, P. S., Donmall, M. C., Mcelduff, P., et al (2006) The role of the early therapeutic alliance in predicting drug treatment dropout. *Drug and Alcohol Dependence*, **83**, 57–64.
- Melnikow, J., Paliescheskey, M. & Stewart, G. K. (1997) Effect of a transportation incentive on compliance with the first prenatal appointment: a randomized trial. *Obstetrics and Gynecology*, **89**, 1023–1027.
- Melo A. P. S. & Guimarães, M. D. C. (2005) Factors associated with psychiatric treatment dropout in a mental health reference center, Belo Horizonte (Fatores associados ao abandono do tratamento psiquiátrico em um centro de referência em saúde mental em Belo Horizonte). *Revista Brasileira de Psiquiatria*, **27**, 113–188.
- Mitchell, A. (2007) Adherence behaviour and psychotropic medication: is it a form of self-medication? *Medical Hypotheses*, **68**, 12–21.
- Mitchell, A., Kingdon, D. & Cross, I. (2006) Management of self-harm following hospital discharge: role for general practitioners in continuity of care. *Primary Care and Community Psychiatry*, **10**, 149–158.
- Mitchell, A. J. & Selmes, T. (2007a) Why don't patients take their medicine? Reasons and solutions in psychiatry. *Advances in Psychiatric Treatment*, **13**, 336–346.
- Mitchell, A. J. & Selmes, T. (2007b) A comparative survey of missed initial and follow-up appointments to psychiatric specialties in the United Kingdom. *Psychiatric Services*, **58**, 868–871.
- Moore, P. J., Sickel, A. E., Malat, J., et al (2004) Psychosocial factors in medical and psychological treatment avoidance: the role of the doctor-patient relationship. *Journal of Health Psychology*, **9**, 421–433.
- Nakanishi, M., Koyama, A., Ito, H., et al (2006) Nurses' collaboration with physicians in managing medication improves patient outcome in acute psychiatric care. *Psychiatry and Clinical Neurosciences*, **60**, 196–203.
- Nelson, E. A., Maruish, M. E. & Axler, J. L. (2000) Effects of discharge planning and compliance with outpatient appointments on readmission rates. *Psychiatric Services*, **51**, 885–889.
- Neupert, S. & McDonald-Miszczak, L. (2004) Younger and older adults' delayed recall of medication instructions: the role of cognitive and metacognitive predictors. *Aging Neuropsychology and Cognition*, **11**, 428–442.
- Nose, M., Barbui, C. & Tansella, M. (2003) How often do patients with psychosis fail to adhere to treatment programmes? A systematic review. *Psychological Medicine*, **33**, 1149–1160.
- Ogrodniczuk, J. S., Joyce, A. S. & Piper, W. E. (2005) Strategies for reducing patient-initiated premature termination of psychotherapy. *Harvard Review of Psychiatry*, **13**, 57–70.
- Pang, A. H., Lum, F. C., Ungvari, G. S., et al (1996) A prospective outcome study of patients missing regular psychiatric outpatient appointments. *Social Psychiatry and Psychiatric Epidemiology*, **31**, 299–302.
- Parrish, J., Charlop, M. & Fenton, L. (1986) Use of a stated waiting list contingency and reward opportunity to increase appointment keeping in an outpatient pediatric psychiatric clinic. *Journal of Pediatric Psychology*, **11**, 81–89.
- Patel, N., Crismon, M., Miller, A., et al (2005) Drug adherence: effects of decreased visit frequency on adherence to clozapine therapy. *Pharmacotherapy*, **25**, 1242–1247.
- Pekarik, G. (1983) Improvement in clients who have given different reasons for dropping out of treatment. *Journal of Clinical Psychology*, **39**, 909–913.
- Percudani, M., Belloni, G., Contini, A., et al (2002) Monitoring community psychiatric services in Italy: differences between patients who leave care and those who stay in treatment. *British Journal of Psychiatry*, **180**, 254–259.
- Pesata, V., Pallija, G. & Webb, A. A. (1999) A descriptive study of missed appointments: families' perceptions of barriers to care. *Journal of Pediatric Health*, **13**, 178–182.
- Pettinati, H. M., Weiss, R. D., Miller, W. R., et al (2004) *Medical Management Treatment Manual: A Clinical Research Guide for Medically Trained Clinicians Providing Pharmacotherapy as Part of the Treatment for Alcohol Dependence*. National Institute on Alcohol Abuse and Alcoholism. http://pubs.niaaa.nih.gov/publications/combine/medication_compliance.htm

Pollastri, A. R., Pokrywa, M. L., Walsh, S. J., et al (2005) Incentive program decreases no-shows in nontreatment substance abuse research. *Experimental and Clinical Psychopharmacology*, **13**, 376–380.

Reda, S. & Makhoul, S. (2001) Prompts to encourage appointment attendance for people with serious mental illness. *Cochrane Database of Systematic Reviews*, issue 2. Oxford Update Software. Art. no.: CD002085. DOI: 10.1002/14651858.CD002085.

Rosen, M. I., Beauvais, J. E., Rigsby, M. O., et al (2003) Neuro-psychological correlates of suboptimal adherence to metformin. *Journal of Behavioral Medicine*, **26**, 349–360.

Rossi, A., Amaddeo, F., Bisoffi, G., et al (2002) Dropping out of care: inappropriate terminations of contact with community-based psychiatric services. *British Journal of Psychiatry*, **181**, 331–338.

Royal College of Psychiatrists (1996) *Report of the Confidential Inquiry into Homicides and Suicides by Mentally Ill People*. Royal College of Psychiatrists.

Sainsbury Centre for Mental Health (2001) *Mental Health Topics: Assertive Outreach*. Sainsbury Centre for Mental Health.

Sanz, M., Constable, G., Lopez-Ibor, L., et al (1998) A comparative study of insight scales and their relationship to psychopathological and clinical variables. *Psychological Medicine*, **28**, 437–446.

Sharp, D. & Hamilton, W. (2001) Nonattendance at general practices and outpatient clinics. *BMJ*, **323**, 1081–1082.

Shivack, I. M. & Sullivan, C. W. (1989) Use of telephone prompts at an inner-city outpatient clinic. *Hospital and Community Psychiatry*, **40**, 851–853.

Smoller, J. W., McLean, R. Y., Otto, M. W., et al (1998) How do clinicians respond to patients who miss appointments? *Journal of Clinical Psychiatry*, **59**, 330–338.

Sparr, L. F., Moffitt, M. C. & Ward, M. F. (1993) Missed psychiatric appointments: who returns and who stays away. *American Journal of Psychiatry*, **150**, 801–805.

Stone, C., Palmer, J., Saxby, P., et al (1999) Reducing nonattendance at outpatient clinics. *Journal of the Royal Society of Medicine*, **92**, 114–118.

Tehrani, E., Krussel, J., Borg, L., et al (1996) Dropping out of psychiatric treatment: a prospective study of a first-admission cohort. *Acta Psychiatrica Scandinavica*, **94**, 266–271.

Waller, J. & Hodgkin, P. (2000) Defaulters in general practice: who are they and what can be done about them. *Family Practice*, **17**, 252–253.

Weinerman, R., Glossop, V., Wong, R., et al (2003) Time of day influences nonattendance at urgent short-term mental health unit in Victoria, British Columbia. *Canadian Journal of Psychiatry*, **48**, 342–344.

Wenning, K. & King, S. (1995) Parent orientation meetings to improve attendance and access at a child psychiatric clinic. *Psychiatric Services*, **46**, 831–833.

Wharne, S. (2005) Assertive outreach teams: their roles and functions. *Journal of Interprofessional Care*, **19**, 326–337.

Wilder, J., Plutchnik, R. & Conte, H. (1977) Compliance with psychiatric emergency room referrals. *Archives of General Psychiatry*, **34**, 930–933.

Young, A. S., Grusky, O., Jordan, D., et al (2000) Routine outcome monitoring in a public mental health system: the impact of patients who leave care. *Psychiatric Services*, **51**, 85–91.

- c in most psychiatric specialties, follow-up patients are less likely to attend than those who are being seen for the first time
- d a missed appointment most commonly heralds complete disengagement from mental health services
- e most cases of missed appointments are due to loss of insight.

2 Psychiatric patients who fail to attend an initial clinic appointment:

- a are generally less unwell than those who do attend
- b may be assumed to be of low risk of suicide
- c are more likely to adhere to medication regimens
- d will be more likely to attend in the future if contacted by telephone
- e should be discharged without follow-up, to avoid wasting NHS resources.

3 The following groups have relatively high rates of non-attendance:

- a older people
- b people given a rapid appointment
- c those who score highly for neuroticism on the Eysenck Personality Questionnaire
- d patients who misuse alcohol and drugs
- e people from higher socio-economic groups.

4 Referrals made by the following groups have been associated with consistently good attendance rates:

- a community psychiatric nurses
- b the probation service
- c patients themselves (self-referral)
- d general practitioners
- e professionals who perceive psychiatry negatively.

5 The following potential interventions to increase attendance have no evidence-based support:

- a telephone reminder calls
- b 'orientation statements'
- c assertive outreach services
- d sending three repeat appointments by post
- e memory aids for people with cognitive impairment.

MCQs

1 Regarding non-attendance at psychiatric out-patient clinics:

- a it is typically less serious than in other medical specialties
- b non-attendance following hospital admission predicts subsequent readmission

MCQ answers

1	2	3	4	5
a F	a F	a F	a T	a F
b T	b F	b F	b F	b F
c F	c F	c F	c F	c F
d F	d T	d T	d F	d T
e F	e F	e F	e F	e F

BJPsych

Advances

Why don't patients attend their appointments? Maintaining engagement with psychiatric services

Alex J. Mitchell and Thomas Selmes

APT 2007, 13:423-434.

Access the most recent version at DOI: [10.1192/apt.bp.106.003202](https://doi.org/10.1192/apt.bp.106.003202)

References

This article cites 85 articles, 14 of which you can access for free at:
<http://apt.rcpsych.org/content/13/6/423#BIBL>

Reprints/ permissions

To obtain reprints or permission to reproduce material from this paper, please write to permissions@rcpsych.ac.uk

You can respond to this article at

[/letters/submit/aptrcpsych;13/6/423](http://apt.rcpsych.org/letters/submit/aptrcpsych;13/6/423)

Downloaded from

<http://apt.rcpsych.org/> on January 20, 2018
Published by [The Royal College of Psychiatrists](http://www.rcpsych.ac.uk/)

To subscribe to *BJPsych Advances* go to:
<http://apt.rcpsych.org/site/subscriptions/>
